

## Cochlear implantation in atelectasis and chronic otitis media: long-term follow-up.

Written by Xenelis Ioannis, Prof. of Otorhinolaryngology - Last Updated Saturday, 06 October 2012 20:09

---



**Xenelis Ioannis, MD** , Professor of Otorhinolaryngology, ENT Department, Athens University, Hipp

**OBJECTIVE:** To report the long-term results of cochlear implantation in cases with chronic otitis media or atelectasis using a single surgical technique performed in a single cochlear implant center.

**PATIENTS:** Nine patients who were implanted using the blind-pit closure of the external ear canal technique (4 patients with adhesive otitis media and 5 with radical mastoid cavities). Follow-up ranged from 18 months to 12 years (mean, 7.05 yr).

**INTERVENTION:** The surgical procedure was performed in 2 stages. The first stage included canal wall down or lowering any high facial ridge in previous mastoidectomies, removal of all skin, and blind-pit closure of the external ear canal without mastoid cavity obliteration or eustachian tube obliteration. Cochlear implantation was performed 6 months after the first surgical procedure.

**RESULTS:** All operations were uneventful, and during cochlear implantation, as a second stage, no epithelia or other problems were encountered. No serious complications were encountered during the follow-up period. One case had a minor disruption of the external canal closure that was reclosed successfully under local anesthesia. All patients were using the device at the last follow-up interval with no device problems.

**CONCLUSION:** Blind-sac closure of the external ear canal without obliteration is a rather safe surgical procedure in cases with chronic otitis media or atelectasis. Meticulous surgical technique and proper patient selection are of paramount importance. However, a 2-stage

## **Cochlear implantation in atelectasis and chronic otitis media: long-term follow-up.**

Written by Xenelis Ioannis, Prof. of Otorhinolaryngology - Last Updated Saturday, 06 October 2012 20:09

---

procedure may not always be necessary and might best be confined to those patients who have active inflammatory disease at the primary procedure.

[Xenellis J](#) , [Nikolopoulos TP](#) , [Marangoudakis P](#) , [Vlastarakos PV](#) , [Tsangaroulakis A](#) , [Ferekidis E](#)

. Cochlear implantation in atelectasis and chronic otitis media: long-term follow-up. Otol Neurotol. 2008 Jun;29(4):499-501